arXiv:math/0410196v1 [math.DG] 7 Oct 2004

RIGIDITY OF SINGULAR SCHUBERT VARIETIES IN $Gr(m,n)$

JAEHYUN HONG

Abstract. Let $\mathbf{a} = (p_1^{q_1}, \cdots, p_r^{q_r})$ be a partition and $\mathbf{a}' = ({p'_1}^{q'_1}, \cdots, {p'_r}^{q'_r})$ be its conjugate. We will prove that if $q_i, q_i \geq 2$ for all $1 \leq i \leq r$, then any irreducible subvariety X of $Gr(m,n)$ whose homology class is an integral multiple of the Schubert class $[\sigma_{\mathbf{a}}]$ of type $\mathbf{a}$ is a Schubert variety of type $\mathbf{a}$.

1. Introduction

Let $Gr(m,n)$ be the Grassmannian of m-planes in $\mathbb{C}^n$. For a partition $\mathbf{a} = (a_1, \cdots, a_m)$ a Schubert variety $\sigma_{\mathbf{a}}$ of type $\mathbf{a}$ is defined by the set of all m-planes E such that $\dim (E \cap \mathbb{C}^{n-m+i-a_i}) \geq i, 1 \leq i \leq m$ for a choice of a flag $\{\mathbb{C}^1 \subset \cdots \subset \mathbb{C}^n\}$. Then they form a basis for the homology space $H_*(Gr(m,n), \mathbb{Z})$.

For $\mathbf{a} = (p^q)^* = ((n-m)^{m-q}, (n-m-p)^q)$, the Schubert variety $\sigma_{\mathbf{a}}$ of type $\mathbf{a}$ is smooth and they are all the smooth Schubert varieties in $Gr(m,n)$. The *Schur rigidity* of smooth Schubert varieties in $Gr(m,n)$ is proved in [W], [B] and [Ho]: for any smooth Schubert variety $\sigma_{\mathbf{a}}$ in $Gr(m,n)$ other than a non-maximal linear space, any irreducible subvariety whose homology class is an integral multiple of the Schubert class $[\sigma_{\mathbf{a}}]$ of type $\mathbf{a}$ is a Schubert variety of type $\mathbf{a}$.

In this paper we will prove the Schur rigidity of singular Schubert varieties of certain types in $Gr(m,n)$.

Theorem. *Let* $\mathbf{a} = (p_1^{q_1}, \cdots, p_r^{q_r})$ *be a partition and let* $\mathbf{a}' = ({p'_1}^{q'_1}, \cdots, {p'_s}^{q'_r})$ *be its conjugate. Then the Schubert variety* $\sigma_{\mathbf{a}}$ *is Schur rigid if* $q_i, q'_i \geq 2$ *for all* $i \leq r$.

The proof is divided by two parts as in [W], [B] and [Ho]: Schubert rigidity and the equality $\mathcal{B}_{\mathbf{a}} = \mathcal{R}_{\mathbf{a}^*}$. Define the Schubert differential system $\mathcal{B}_{\mathbf{a}}$ by the differential system with a fiber at $x \in Gr(m,n)$ given by the set of all the tangent space of Schubert varieties of type $\mathbf{a}$ passing through x. If any irreducible integral variety of $\mathcal{B}_{\mathbf{a}}$ is a Schubert variety of type $\mathbf{a}$, then we say that the Schubert variety $\sigma_{\mathbf{a}}$ is *Schubert rigid*.

Putting together the tangent space of all the subvarieties X with $[X] = r[\sigma_{\mathbf{a}}]$, $r \in \mathbb{Z}$ at each point, we get another differential system $\mathcal{R}_{\mathbf{a}^*}$, which we call the Schur differential system([B], [W]). By the construction, if any irreducible integral variety of $\mathcal{R}_{\mathbf{a}^*}$ is a Schubert variety of type $\mathbf{a}$, then the Schubert variety $\sigma_{\mathbf{a}}$ is Schur rigid. Furthermore, the Schubert differential system $\mathcal{B}_{\mathbf{a}}$ is always contained in the Schur differential system $\mathcal{R}_{\mathbf{a}^*}$. Thus the equality $\mathcal{B}_{\mathbf{a}} = \mathcal{R}_{\mathbf{a}^*}$ and the Schubert rigidity is a necessary and sufficient condition for the Schur rigidity([B]). While proving the equality $\mathcal{B}_{\mathbf{a}} = \mathcal{R}_{\mathbf{a}^*}$ is computing integral elements of exterior differential systems, which is an algebraic problem, proving the Schubert rigidity is finding integral varieties of a differential system, which is a local differential geometric problem.

When the Schubert variety $\sigma_{\mathbf{a}}$ is singular, we will express the Schubert variety $\sigma_{\mathbf{a}}$ as the locus of a family of smooth Schubert varieties $\sigma_{\mathbf{b}}$ of type $\mathbf{b}$ on $Gr(m,n)$ parameterized by a Schubert variety $\sigma_{\mathbf{c}}$ of type $\mathbf{c}$ in another Grassmannian which is the parameter space of the Schubert variety of type $\mathbf{b}$ on $Gr(m,n)$. Then the Schubert variety $\sigma_{\mathbf{a}}$ is Schubert rigid if the Schubert varietyies $\sigma_{\mathbf{b}}$ and $\sigma_{\mathbf{c}}$ are Schubert rigid. So we can use the Schubert rigidity of smooth Schubert varieties and the induction to prove the Schubert rigidity of singular Schubert varieties in $Gr(m,n)$ of types in Theorem(Section 3).

To prove the equality $B_{\mathbf{a}} = R_{\mathbf{a}^*} \subset Gr(k, E^* \otimes Q)$ for the fibers of $\mathcal{B}_{\mathbf{a}}$ and $\mathcal{R}_{\mathbf{a}^*}$ at $[E] \in Gr(m,n)$, we use the description $R_{\mathbf{a}} = Gr(|\mathbf{a}|, E^* \otimes Q) \cap \mathbb{P}(\mathbb{S}_{\mathbf{a}}(E^*) \otimes \mathbb{S}_{\mathbf{a}'}(Q))$ as in [Ho] and then we compute the complement of the tangent space of $B_{\mathbf{a}}$ in the tangent space of $Gr(k, E^* \otimes Q)$ by hands, while in [Ho] the theory of Lie algebra cohomology developed by Kostant used to compute it(Section 4).

Acknowledgements I wish to thank Professor D. Burns for suggesting the idea of using a natural foliation by smooth Schubert varieties on singular Schubert varieties to prove their rigidity. Some results of the present paper have been worked out while I was visiting MSRI in Berkeley in November 2003. I would like to thank this institute for its hospitality, and Professor R. Bryant, one of the organizers of the program “Differential Geometry”, for stimulating discussions and encouragement. This work was supported by the Post-Doctorial Fellowship Program of Korea Science and Engineering Foundation(KOSEF).

2. DIFFERENTIAL SYSTEMS

2.1. **Schubert differential systems.** Let $Gr(m,n)$ be the Grassmannian of m-dimensional subspaces of $V = \mathbb{C}^n$. Let $P(m,n)$ be the set

$$\{\mathbf{a} = (a_1, \cdots, a_m) |\ n-m \geq a_1 \geq \cdots \geq a_m \geq 0\}$$

of partitions. Fix a flag $\{V_\bullet\}$ of V with $\dim V_i = i$. For $\mathbf{a} \in P(m,n)$, define the *Schubert variety* $\sigma_{\mathbf{a}}(V_\bullet)$ *of type* $\mathbf{a}$ by the set

$$\{E \in Gr(m,n) |\ \dim(E \cap V_{n-m+i-a_i}) \geq i\}.$$

Then $\sigma_{\mathbf{a}}(V_\bullet)$ is a subvariety of $Gr(m,n)$ of codimension $|\mathbf{a}| := a_1 + \cdots + a_m$. By varying the flag $\{V_\bullet\}$, we get a family of Schubert varieties $\sigma_{\mathbf{a}}$ of type $\mathbf{a}$.

Let $\mathbf{a} \in P(m,n)$. Define its *dual* $\mathbf{a}^*$ by

$$\mathbf{a}^* = (n-m-a_m, \cdots, n-m-a_1)$$

and define its *conjugate* $\mathbf{a}' = (a'_1, \cdots, a'_{n-m})$ by

$$a'_i = \sharp\{j | a_j \geq i\} \text{ for } 1 \leq i \leq n-m.$$

The *Young diagram* $Y_{\mathbf{a}}$ is defined by the set of boxes consisting of a_i boxes in the i-th row, the row of boxes lined up on the left. Then the Young diagram $Y_{\mathbf{a}^*}$ is obtained by rotating the complement of the Young diagram $Y_{\mathbf{a}}$ by 180 degree and the Young diagram $Y_{\mathbf{a}'}$ is obtained by transposing the Young diagram $Y_{\mathbf{a}}$.

Example 2.1. $m = 5$, $n - m = 6$. Young diagrams $Y_{\mathbf{a}}, Y_{\mathbf{a}^*}, Y_{\mathbf{a}'}$ for $\mathbf{a} = (6,6,4,2,2)$. $\mathbf{a}^* = (4,4,2)$, $\mathbf{a}' = (5,5,3,3,2,2)$







Define $\mathfrak{n}_{\mathbf{a}}$ to be the vector space of matrices $Z = (z^a_i) \in M_{n-m,m}$ that satisfy $z^a_i = 0$ when $a > n-m-a_i$. This is the tangent space of the Schubert variety $\sigma_{\mathbf{a}}$ and is obtained by rotating the Young diagram $Y_{\mathbf{a}^*}$ by 90 degree in counterclockwise.



We will use the notation $\mathbf{a} = (p_1^{q_1}, \cdots, p_r^{q_r})$, $p_r \neq 0$ for the partition with q_1 p_1's, $\cdots$, q_r p_r's.

Definition 2.2. Let M be a manifold and let $Gr(k, TM)$ be the Grassmannian bundle of k-subspaces of the tangent bundle TM. A subvariety $\mathcal{F}$ of $Gr(k, TM)$ is called a *differential system* on M. A subvariety X of M is said to be an *integral variety* of the differential system $\mathcal{F}$ if at each smooth point $x \in X$, the tangent space T_xX is an element of the fiber $\mathcal{F}_x$. We say that $\mathcal{F}$ is *integrable* if at each point $x \in M$ and $y \in \mathcal{F}_x$, there is an integral variety passing through x and tangent to the subspace W_y of T_xM corresponding to y.

Definition 2.3. For each $\mathbf{a} \in P(m,n)$, the *Schubert differential system* $\mathcal{B}_{\mathbf{a}}$ of type $\mathbf{a}$ is the differential system with a fiber consisting of the tangent space to the Schubert varieties $\sigma_{\mathbf{a}}$ of type $\mathbf{a}$ passing through a given point. We say that $\sigma_{\mathbf{a}}$ is *Schubert rigid* if Schubert varieties of type $\mathbf{a}$ are the only irreducible integral varieties of $\mathcal{B}_{\mathbf{a}}$. If, furthermore, there is a unique Schubert variety passing through a given point and tangent to a given tangent subspace, then we say that $\mathcal{B}_{\mathbf{a}}$ is *strongly rigid*.

Remark 2.4. For a smooth Schubert variety in $Gr(m,n)$, the Schubert differential system is strongly rigid ([W], Proposition 3.3) and this will play a central role in proving the Schubert rigidity of singular Schubert varieties.

Let $[E] \in Gr(m,n)$ and $Q = \mathbb{C}^n/E$. Then $T_{[E]}Gr(m,n) = E^* \otimes Q$ and $\wedge^k(E^* \otimes Q) = \oplus_{|\mathbf{a}|=k}\mathbb{S}_{\mathbf{a}}(E^*) \otimes \mathbb{S}_{\mathbf{a}'}(Q)$, where $\mathbb{S}_{\mathbf{a}}$ is the Schur functor of type $\mathbf{a}$. There exists a $SU(n)$-invariant positive (k,k)-form $\phi_{\mathbf{a}}$ which can be written as the sum $(\sqrt{-1})^{k^2}\sum_i \xi_i \wedge \overline{\xi}_i$ at $[E]$, where $\{\xi_i\}$ is an orthonormal basis of $\mathbb{S}_{\mathbf{a}}(E) \otimes \mathbb{S}_{\mathbf{a}'}(Q^*) \subset \wedge^k(T^*_{[E]}Gr(m,n))$. Then we have $\int_{\sigma_{\mathbf{a}^*}} \phi_{\mathbf{b}} = \delta^{\mathbf{b}}_{\mathbf{a}}$ for $\mathbf{a}, \mathbf{b} \in P(m,n)$(For details, see [B]).

Definition 2.5. For each $\mathbf{a} \in P(m,n)$, the *Schur differential system* $\mathcal{R}_{\mathbf{a}}$ of type $\mathbf{a}$ is defined by the intersection

$$\cap_{\mathbf{b}\neq\mathbf{a},|\mathbf{b}|=|\mathbf{a}|}Z(\phi_{\mathbf{b}}),$$

where $Z(\phi_{\mathbf{b}})$ is the set of $|b|$-subspace of $T_E(Gr(m,n))$ on which $\phi_{\mathbf{b}}$ vanishes. We say that $\sigma_{\mathbf{a}}$ is *Schur rigid* if Schubert varieties of type $\mathbf{a}^*$ are the only integral varieties of $\mathcal{R}_{\mathbf{a}}$.

Remark 2.6. (1) $\mathcal{B}_{\mathbf{a}^*}$ is contained in $\mathcal{R}_{\mathbf{a}}$, and X is a subvariety of $Gr(m,n)$ with $[X] = r[\sigma_{\mathbf{a}^*}]$ for an integer r if and only if X is an integral variety of $\mathcal{R}_{\mathbf{a}}$([B], [W]).

(2) $\mathcal{B}_{\mathbf{a}^*}$ is closed and $\mathcal{R}_{\mathbf{a}}$ is connected (Remark 2 and Remark 12 of [B]).

2.2. F-structures and integral varieties.

Definition 2.7. Let F be a submanifold of $Gr(k,V)$ with a transitive action of a subgroup of $GL(V)$. A fiber bundle $\mathcal{F} \subset Gr(k,TM)$ on a manifold M of dimension $n = \dim V$ is said to be an *F-structure* if at each point $x \in M$ there is a linear isomorphism $\varphi(x) : V \to T_xM$ such that the induced map $\varphi(x)^k : Gr(k,V) \to Gr(k,T_xM)$ sends F to $\mathcal{F}_x$.

Schubert differential systems are F-structures for various F's. Integrability and the uniqueness can be obtained by studying the cohomology space $H^{k,1}(F)$ and $H^{k,2}(F)$ associated to (W_f, V, T_fF) for $f \in F$([G]). When an F-structure is integrable, if $H^{k,1}(F) = 0$, then the family of all integral varieties passing through a fixed point and tangent to a fixed subspace has dimension $\sum_{j\leq k-1}\dim H^{j,1}(F)$. In particular, if $H^{1,1}(F) = 0$, there is only one such integral subvariety. Higher cohomology gives the information on the higher jet of the integral varieties.

In general, Schubert differential system for singular Schubert variety has order ≥ 2, i.e. its integral varieties are determined by higher jets. But smooth Schubert varieties depend only 1-jet and there is a canonical map from the differential system of a singular Schubert variety to that of a certain smooth Schubert variety(Proposition 3.2). So, in this paper, we will consider only the first cohomology $H^{1,1}(F)$, which contains the information on the 2-jets of integral varieties.

Definition 2.8. Let F be a subvariety of $Gr(k,V)$ with a transitive action of a subgroup of $GL(V)$. Let $\mathcal{F}$ be an F-structure on M with the projection map $\pi : \mathcal{F} \to M$. For $x \in M$ and $y \in \mathcal{F}_x$, let W_y denote the k-subspace of T_xM corresponding to y. For a k-subspace H of $T_y\mathcal{F}$ such that $\pi_* : H \to W_y$ is an isomorphism, define $\partial H : \wedge^2 W_y \to T_xM/W_y$ by

$$\partial H(V_1, V_2) = [\tilde{V_1}, \tilde{V_2}] \mod W_y,$$

$\tilde{V_i}, i = 1,2$ is a local vector field on M with $\tilde{V_i}(x') \in W_{\psi(x')}$ for a local section ψ of $\mathcal{F}$ with $\psi_*(W_y) = H$. It is well defined([G]).

A k-subspace H of $T_y\mathcal{F}$ is said to be a *2-jet of an integral variety* if π_* restricts to an isomorphism $H \subset T_y\mathcal{F} \to W_y \subset T_xM$ and $\partial H = 0$.

Such an H is indeed a candidate for the 2-jets of actual integral varieties of $\mathcal{F}$ and the set of such an H is again a subvariety of $Gr(k, V + T_fF)$.

Proposition 2.9. *Let $\mathcal{F}$ be an F-structure on M. Let $x \in M$ and let $y \in \mathcal{F}_x$. If X is an integral variety of $\mathcal{F}$ passing through x and tangent to W_y, then the tangent space $H = T_y\tilde{X} \subset T_y\mathcal{F}$ of the lifting $\tilde{X} := \{(x, [T_xX]) | x \in X, [T_xX] \in Gr(k, T_xM)\}$ of X to $Gr(k, T_xM)$ satisfies*

(1) $\pi_* : H \to W_y$ is an isomorphism

(2) $\partial H = 0$.

Proof. See Chapter 1 of [G]. □

The condition that $\pi_* : H \to W_y$ is an isomorphism is equivalent to the condition that H is the graph of a map $p : W_y \to T_y(\mathcal{F}_x) \subset W_y^* \otimes (T_xM/W_y)$. Define $\partial p : \wedge^2 W_y \to T_xM/W_y$ by $\partial p(V_1, V_2) = p(V_1)(V_2) - p(V_2)(V_1)$, considering $T_y(\mathcal{F}_x)$ as a subspace of $W_y^* \otimes (T_xM/W_y)$. Then $\partial H = 0$ if and only if $\partial p = 0$.

Definition 2.10. Let $\mathcal{F}$ be an F-structure on M. For each $y \in \mathcal{F}$, define $\mathcal{F}_y^{(1)}$ by the set of all 2-jets of integral varieties tangent to W_y, i. e.,

$$\mathcal{F}_y^{(1)} = \{H \subset T_y\mathcal{F} : \pi_* : H \to W_y \text{ is an isomorphism and } \partial H = 0\}.$$

We call $\mathcal{F}^{(1)} = \cup_{y \in \mathcal{F}} \mathcal{F}_y^{(1)}$ the *first prolongation* of $\mathcal{F}$. Define

$$F^{(1)} = \{H \subset V + T_fF | H \text{ is the graph of a map } p : W_f \to T_fF,\ \partial p = 0\}.$$

Then $\mathcal{F}^{(1)}$ is an $F^{(1)}$-structure on $\mathcal{F}$.

Put $H^{1,1}(F) = Ker(\partial : W_f^* \otimes T_f F \to \wedge^2 W_f^* \otimes (V/W_f))$. If $H^{1,1}(F) = 0$ then $F^{(1)}$ is just a point and the first prolongation defines a distribution on $\mathcal{F}$. In this case, the integrability of this distribution is equivalent to the integrability of the F-structure $\mathcal{F}$. So there is at most one integral manifold passing through a given point and tangent to a given k-subspace of the tangent space. For the details see Chapter 1 of [G].

3. Schubert rigidity

3.1. **Description of the Schubert differential system.**

Example 3.1. For $\mathbf{a} = (p^q)^*$, The Schubert variety $\sigma_{\mathbf{a}}$ is the sub-Grassmannian $\{E \in Gr(m,n) | \mathbb{C}^{m-q} \subset E \subset \mathbb{C}^{m+p}\} \simeq Gr(q, p+q)$. The Schubert differential system $\mathcal{B}_{\mathbf{a}}$ is the flag space $F(m-q, m, m+p, n)$ and the parameter space of the family of Schubert varieties of type $\mathbf{a}$ is $F(m-q, m+p, n)$. In this case, there is a double fibration $F(m-q, m+p, n) \leftarrow \mathcal{B}_{\mathbf{a}} \to Gr(m,n)$.

In general, the Schubert differential systems $\mathcal{B}_{\mathbf{a}}$ is a generalized flag variety. Since $SL(n)$ acts on $\mathcal{B}_{\mathbf{a}}$ transitively and $\mathcal{B}_{\mathbf{a}}$ is compact, we have only to find the corresponding subset of simple root system which generate the isotropy group which is parabolic.

Let $\mathcal{S} = \{\alpha_1, \cdots, \alpha_{n-1}\}$ be the set of simple roots of $G = SL(n)$ and let P be the parabolic subgroup of $SL(n)$ generated by $\mathcal{S}^1 = \mathcal{S} - \{\alpha_m\}$, i.e. $\mathcal{S}^1$ is the set of simple roots of the semisimple part $SL(m) \times SL(n-m)$ of P. Then $Gr(m,n)$ is G/P.

Proposition 3.2. *For* $\mathbf{a} = (p_1^{q_1}, \cdots, p_r^{q_r}), p_r \neq 0$, *put*

$$\mathcal{S}_{\mathbf{a}} = \mathcal{S}^1 - \{\alpha_{q_1}, \cdots, \alpha_{q_1+\cdots+q_r}\} \cup \{\alpha_{n-p_1} \cdots, \alpha_{n-p_r}\}.$$

Let $Q_{\mathbf{a}}$ *is the parabolic subgroup of* $SL(n)$ *generated by* $\mathcal{S}_{\mathbf{a}} \cup \{\alpha_m\}$ *and* $P_{\mathbf{a}}$ *be the parabolic subgroup of* $SL(m) \times SL(n-m)$ *generated by the set* $\mathcal{S}_{\mathbf{a}}$. *Then* $\mathcal{B}_{\mathbf{a}}$ *is the homogeneous manifold* $G/(Q_{\mathbf{a}} \cap P)$ *and* $P_{\mathbf{a}}$ *is the isotropy group of the the action of* $SL(m) \times SL(n-m)$ *on the fiber* $B_{\mathbf{a}}$ *of* $\mathcal{B}_{\mathbf{a}} \to Gr(m,n)$. *If* $\mathcal{S}_{\mathbf{a}} \subset \mathcal{S}_{\mathbf{b}}$, *then there exists a quotient map* $\varphi_{\mathbf{a},\mathbf{b}} : \mathcal{B}_{\mathbf{a}} \to \mathcal{B}_{\mathbf{b}}$ *which preserves the fibers.*

For example, the following figure describes $\mathfrak{n}_{\mathbf{a}}$, $\mathfrak{m}_{\mathbf{a}}$ and $Q_{\mathbf{a}}$ for $n = 10, m = 4$ and $\mathbf{a} = (6,4,2,2)$, where $\mathfrak{m}_{\mathbf{a}}$ denotes the tangent space of $B_{\mathbf{a}}$. $\mathfrak{n}_{\mathbf{a}}$ is the space of all $n \times n$-matrix with nonzero elements only in $*$ and $\mathfrak{m}_{\mathbf{a}}$ is the space of all $n \times n$-matrix with nonzero elements only in $\bullet$. The Lie algebra of the reductive part of $Q_{\mathbf{a}}$ is the space of all $n \times n$-matrix with nonzero elements only in $\diamond$.

◇									
●	◇								
●	●	◇	◇						
●	●	◇	◇						
	*	*	*	◇	◇				
	*	*	*	◇	◇				
		*	*	●	●	◇	◇		
		*	*	●	●	◇	◇		
				●	●	●	●	◇	◇
				●	●	●	●	◇	◇

Proposition 3.3. *For* $\mathbf{a} = (p^q)^*$*, the Schubert variety* $\sigma_{\mathbf{a}}$ *of type* $\mathbf{a}$ *is strongly rigid except when* ($p = 1$ *and* $q \neq m$) *or* ($p \neq n - m$ *and* $q = 1$)*.*

Proof. See [W]. □

3.2. **Foliation by smooth Schubert varieties.** We start with the simplest case and then will use the induction to prove the general case.

Example 3.4. The case when $\mathbf{a} = (p^q)$ is studied in Example 13 and Remark 33 of [B]. Fix a $(n-m-p+q)$-subspace $\Lambda = \mathbb{C}^{n-m-p+q}$ of $\mathbb{C}^n$. Then $\sigma_{\mathbf{a}}(\Lambda)$ can be expressed as a union of a family of Schubert varieties of type $\mathbf{b} = ((n-m)^q) = ((n-m)^{(m-q)})^*$.

$$\begin{aligned}\sigma_{\mathbf{a}}(\Lambda) &= \{E \in Gr(m,n) | \dim(E \cap \Lambda) \geq q\} \\ &= \bigcup_{\mathbb{C}^q \subset \Lambda} \{E \in Gr(m,n) | \mathbb{C}^q \subset E\}\end{aligned}$$

Note that $Gr(q, n)$ is the parameter space of the Schubert varieties of type $\mathbf{b}$ and $\{\mathbb{C}^q \in Gr(q,n) | \mathbb{C}^q \subset \Lambda\}$ is a Schubert variety of type $\mathbf{c} = ((n-m-p)^q)^*$ in $Gr(q, n)$.

$$\mathbf{a} = \begin{array}{|c|c||c|c|}\hline 0 & 0 & 0 & 0 \\ \hline \mathfrak{m}_{\mathbf{a}} & 0 & 0 & 0 \\ \hline\hline \mathfrak{n}_{\mathbf{a}} & \mathfrak{n}_{\mathbf{a}} & 0 & 0 \\ \hline 0 & \mathfrak{n}_{\mathbf{a}} & \mathfrak{m}_{\mathbf{a}} & 0 \\ \hline\end{array} \qquad \mathbf{b} = \begin{array}{|c|c||c|c|}\hline 0 & 0 & 0 & 0 \\ \hline \mathfrak{m}_{\mathbf{b}} & 0 & 0 & 0 \\ \hline\hline 0 & \mathfrak{n}_{\mathbf{b}} & 0 & 0 \\ \hline 0 & \mathfrak{n}_{\mathbf{b}} & 0 & 0 \\ \hline\end{array} \qquad \mathbf{c} = \begin{array}{|c||c|c|c|}\hline 0 & 0 & 0 & 0 \\ \hline\hline 0 & \mathfrak{m}_{\mathbf{c}} & 0 & 0 \\ \hline \mathfrak{n}_{\mathbf{c}} & 0 & 0 & 0 \\ \hline 0 & \mathfrak{m}_{\mathbf{c}} & 0 & 0 \\ \hline\end{array}$$

From this expression, we get the following desingularisation π_2 of $\sigma_{\mathbf{a}}(\Lambda)$.

$$\begin{tikzcd} \{(\mathbb{C}^q, E) | \mathbb{C}^q \subset E\} \arrow[r, "\pi_2"] \arrow[d, "\pi_1"'] & Gr(m,n) \supset \sigma_{\mathbf{a}}(\Lambda) \\ Gr(q,n) \supset Gr(q,\Lambda) & \end{tikzcd}$$

Here, $\sigma_{\mathbf{a}}(\Lambda)$ is equal to $\pi_2(\pi_1^{-1}(Gr(q,\Lambda)))$ and $\pi_2 : \pi_1^{-1}(Gr(q,\Lambda)) \to \sigma_{\mathbf{a}}(\Lambda)$ is generically one-to-one.

The smooth locus of $\sigma_{\mathbf{a}}(\Lambda)$ is foliated by Schubert varieties of type $\mathbf{b}$ in $Gr(m,n)$ and the space of leaves of this foliation is a Schubert variety of type $\mathbf{c}$ in $Gr(q,n)$.

In the same way as above, we get a different desingularisation of $\sigma_{\mathbf{a}}(\Lambda)$ by considering $\sigma_{\mathbf{a}}(\Lambda)$ as a union of a family of another type of Schubert varieties:

$$\sigma_{\mathbf{a}}(\Lambda) = \bigcup_{\Lambda \subset \mathbb{C}^{n-p}} \{E \in Gr(m,n) | E \subset \mathbb{C}^{n-p}\}.$$

Proposition 3.5. *For the partition $\mathbf{a} = (p^q)$, the Schubert variety $\sigma_{\mathbf{a}}$ of type $\mathbf{a}$ is rigid if $p > 1$ and $q > 1$.*

It is proved in Example 13 and Remark 33 of [B]. We will prove it again in such a way that can be generalized to the case of the Schubert differential systems of other singular Schubert varieties.

Lemma 3.6. *Let $\mathbf{a}$ and $\mathbf{b}$ be two partitions such that $\mathfrak{n}_{\mathbf{b}}$ is a subspace of $\mathfrak{n}_{\mathbf{a}}$ with $\mathcal{S}_{\mathbf{a}} \subset \mathcal{S}_{\mathbf{b}}$ (and thus there is a projection $\varphi_{\mathbf{a},\mathbf{b}} : \mathcal{B}_{\mathbf{a}} \to \mathcal{B}_{\mathbf{b}}$ as in proposition 3.2). Assume that $\mathcal{B}_{\mathbf{b}}$ is strongly rigid. For $p : \mathfrak{n}_{\mathbf{a}} \to \mathfrak{m}_{\mathbf{a}}$, define $\tilde{p}$ by the composition $\mathfrak{n}_{\mathbf{a}} \to \mathfrak{m}_{\mathbf{a}} \twoheadrightarrow \mathfrak{m}_{\mathbf{b}}$. If $\tilde{p}(v) = 0$ for all $v \in \mathfrak{n}_{\mathbf{b}}$ and for all $p \in H^{1,1}(B_{\mathbf{a}})$, then the smooth locus of any integral variety of $\mathcal{B}_{\mathbf{a}}$ is foliated by integral varieties of $\mathcal{B}_{\mathbf{b}}$, which are Schubert varieties of type $\mathbf{b}$.*

Proof. By the strong rigidity of $\mathcal{B}_{\mathbf{b}}$, its first prolongation $\mathcal{B}_{\mathbf{b}}^{(1)}$ gives a distribution D on $\mathcal{B}_{\mathbf{b}}$ which is integrable because $\mathcal{B}_{\mathbf{b}}$ is integrable. Integral varieties of D are isomorphic to Schubert varieties of type $\mathbf{b}$ via the map $\pi_{\mathbf{b}} : \mathcal{B}_{\mathbf{b}} \to Gr(m,n)$.

Let X be an integral variety of $\mathcal{B}_{\mathbf{a}}$. Let $\tilde{X} \subset \mathcal{B}_{\mathbf{a}}$ be the lifting of X, that is, $\tilde{X} = \{(x, [T_x X]) | x \in X\}$. It suffices to show that $\varphi_{\mathbf{a},\mathbf{b}}(\tilde{X})$ is foliated by the integral varieties of the distribution D induced by $\mathcal{B}_{\mathbf{b}}^{(1)}$, that is, at each point $y \in \tilde{X}$, $(\varphi_{\mathbf{a},\mathbf{b}})_*(T_y \tilde{X})$ contains $D_{\varphi_{\mathbf{a},\mathbf{b}}(y)}$.

$$\begin{tikzcd}
 & & \mathcal{B}_{\mathbf{b}}^{(1)} \arrow[d] \\
\tilde{X} \subset \mathcal{B}_{\mathbf{a}} \arrow[rr, "\varphi_{\mathbf{a},\mathbf{b}}"] \arrow[dr] & & \mathcal{B}_{\mathbf{b}} \supset \varphi_{\mathbf{a},\mathbf{b}}(\tilde{X}) \arrow[dl] \\
 & X \subset Gr(m,n) &
\end{tikzcd}$$

The map $(\varphi_{\mathbf{a},\mathbf{b}})_*$ is given by the projection $\mathfrak{m} + \mathfrak{m}_{\mathbf{a}} \to \mathfrak{m} + \mathfrak{m}_{\mathbf{b}}$ and $T_y \tilde{X}$ is the graph of a map $p : \mathfrak{n}_{\mathbf{a}} \to \mathfrak{m}_{\mathbf{a}}$ with $\partial p = 0$. So $(\varphi_{\mathbf{a},\mathbf{b}})_*(T_y \tilde{X})$ is the graph of the map $\mathfrak{n}_{\mathbf{a}} \xrightarrow{p} \mathfrak{m}_{\mathbf{a}} \to \mathfrak{m}_{\mathbf{b}}$ with $\partial p = 0$. On the other hand $D_{\varphi_{\mathbf{a},\mathbf{b}}(y)}$ is the graph of the zero map $\mathfrak{n}_{\mathbf{b}} \to \mathfrak{m}_{\mathbf{b}}$. By the assumption, $\tilde{p}(v)$ is zero for all $v \in \mathfrak{n}_{\mathbf{b}}$, so $\varphi_{\mathbf{a},\mathbf{b}}(T_y \tilde{X})$ contains $D_{\varphi_{\mathbf{a},\mathbf{b}}(y)}$. $\square$

Lemma 3.7. *Let* $\mathbf{a}, \mathbf{c}$ *be as in Example 3.4. Let* $A \subset Gr(q,n)$ *be a subvariety of dimension* $q(n-m-p)$*. Define* $X_A := \pi_2(\pi_1^{-1}(A)) \subset Gr(m,n)$*. If* X_A *is an integral variety of* $\mathcal{B}_{\mathbf{a}}(Gr(m,n))$ *and* $dim(\pi_1^{-1}(A))$ *is equal to* $dim(X_A)$*, then* A *is an integral variety of* $\mathcal{B}_{\mathbf{c}}(Gr(q,n))$*.*

$$\begin{tikzcd}
\mathcal{B}_{\mathbf{a}} \arrow[rr] \arrow[dr] & & \mathcal{B}_{\mathbf{b}} \arrow[r, "\pi_1"] \arrow[dl, "\pi_2"] & Gr(q,n) \supset A \\
& Gr(m,n) \supset X_A & &
\end{tikzcd}$$

Proof. We will follow the arguments in Example 16.6 of [H]. Define $\tilde{\pi}_i, i = 1, 2$ to be the projection from $Gr(q,n) \times Gr(m,n)$ to the first and second component, respectively. Let $F(q,m,n)$ denote the flag space $\{(\Gamma, E) | \Gamma \subset E \subset \mathbb{C}^n\} \subset Gr(q,n) \times Gr(m,n)$. Then $\pi_2(\pi_1^{-1}(A))$ is $\tilde{\pi}_2(\tilde{\pi}_1^{-1}(A) \cap F(q,m,n))$.

Let Γ be a smooth point in A. Then $\tilde{\pi}_1^{-1}(A)$ is smooth at (Γ, E) for all $E \in Gr(m,n)$ with the tangent space

$$T_{(\Gamma,E)}\tilde{\pi}_1^{-1}(A) = \left\{ (\eta, \varphi) | \begin{array}{ll} \eta : \Gamma \to \mathbb{C}^n/\Gamma, & \eta \in T_\Gamma A \\ \varphi : E \to \mathbb{C}^n/E, & \end{array} \right\}.$$

The tangent space of $F(q,m,n)$ at (Γ, E) is

$$T_{(\Gamma,E)}F(q,m,n) = \left\{ (\eta, \varphi) | \begin{array}{ll} \eta : \Gamma \to \mathbb{C}^n/\Gamma, & \\ \varphi : E \to \mathbb{C}^n/E, & \varphi|_\Gamma \equiv \eta \mod E \end{array} \right\}.$$

By dimension counting, we see that the two tangent spaces are transversal so that $\pi_1^{-1}(A) = \tilde{\pi}_1^{-1}(A) \cap F(q,m,n)$ is smooth at all (Γ, E) with $\Gamma \subset E$, and the tangent space of $\pi_1^{-1}(A)$ at (Γ, E) is given by

$$T_{(\Gamma,E)}\pi_1^{-1}(A) = \left\{ (\eta, \varphi) | \begin{array}{ll} \eta : \Gamma \to \mathbb{C}^n/\Gamma, & \eta \in T_\Gamma A \\ \varphi : E \to \mathbb{C}^n/E, & \varphi|_\Gamma \equiv \eta \mod E \end{array} \right\}.$$

For each $E \in X_A$ there are only finitely many $\Gamma \in A$ with $\Gamma \subset E$. If there are more than one $\Gamma \in A$ with $\Gamma \subset E$, then X_A is not smooth at E(Proposition 16.8 of [H]). Let E be an element in X_A such that there is only one $\Gamma \in A$ with $\Gamma \subset E$. Then $\pi_2 : \pi_1^{-1}(A) \to X_A$ is one-to-one over E so X_A is smooth at E with the tangent space

$$T_E(X_A) = \{\varphi : E \to \mathbb{C}^n/E \mid \varphi|_\Gamma \equiv \eta \mod E, \ \eta \in T_\Gamma A, \}.$$

Since $T_E(X_A)$ is of type $\mathbf{a}$ in $Gr(m,n)$, $T_\Gamma A$ is of type $\mathbf{c}$ in $Gr(q,n)$. $\square$

Proof of Proposition 3.5 We will show that for all $p : \mathfrak{n}_{\mathbf{a}} \to \mathfrak{m}_{\mathbf{a}}$ with $p \in H^{1,1}(B_{\mathbf{a}})$, the induced map $\tilde{p} : \mathfrak{n}_{\mathbf{b}} \subset \mathfrak{n}_{\mathbf{a}} \to \mathfrak{m}_{\mathbf{a}} \to \mathfrak{m}_{\mathbf{b}}$ is zero. Then for an integral variety X of $\mathcal{B}_{\mathbf{a}}$, the space of leaves $A = \pi_1(\varphi_{\mathbf{a},\mathbf{b}}(\tilde{X})) \subset Gr(q,n)$ of the foliation on X given by Lemma 3.6 satisfies the conditions in Lemma 3.7. Thus $A \subset Gr(q,n)$ is an integral variety of $\mathcal{B}_{\mathbf{c}}(Gr(q,n))$. Since $q > 1$ $\mathcal{B}_{\mathbf{c}}$ is rigid. So A is the Schubert variety $Gr(q, \Lambda)$ for a $(n-m-p+q)$-subspace Λ of $\mathbb{C}^n$ and hence X is the Schubert variety $\sigma_{\mathbf{a}}(\Lambda)$.

Let $E_{i,j}$ be the $n\times n$-matrix with only one nonzero element in the i-th row and j-th column. Then $[E_{i,j},E_{k,\ell}]=\delta_{j,k}E_{i\ell}-\delta_{i,\ell}E_{k,j}$ for all $1\leq i,j,k,\ell\leq n$.

Since $[p(X),Y]-[p(Y),X]\in\mathfrak{n}_{\mathbf{a}}$ for all $X,Y\in\mathfrak{n}_{\mathbf{a}}$ and $[\mathfrak{m}_{\mathbf{a}}/\mathfrak{m}_{\mathbf{b}},\mathfrak{n}_{\mathbf{b}}]\subset\mathfrak{n}_{\mathbf{b}}$, $[\tilde{p}(X),Y]-[\tilde{p}(Y),X]\in\mathfrak{n}_{\mathbf{a}}$ for all $X,Y\in\mathfrak{n}_{\mathbf{b}}$. Assume that $q+1\leq i,j\leq m$ and $m+1\leq r,s\leq n-p$ and $n-p+1\leq a,b\leq n$. Note that for a fixed a, if $X\in\mathfrak{m}_{\mathbf{b}}$ and $[X,E_{a,i}]=0$ for all i, then $X=0$.

0	0	0	0
$\mathfrak{m}_{\mathbf{b}}$	0	0	0
$*$	$E_{r,i}$	0	0
0	$E_{a,i}$	$\bullet$	0

From $[\tilde{p}(E_{a,i}),E_{r,j}]-[\tilde{p}(E_{r,j}),E_{a,i}]\in\mathfrak{n}_{\mathbf{a}}$, we get both $[\tilde{p}(E_{a,i}),E_{r,j}]$ and $[\tilde{p}(E_{r,j}),E_{a,i}]$ are contained in $\mathfrak{n}_{\mathbf{a}}$. Thus $[\tilde{p}(E_{r,j}),E_{a,i}]$ is zero, which implies that $\tilde{p}(E_{r,j})=0$.

Since $[\tilde{p}(E_{a,i}),E_{b,j}]-[\tilde{p}(E_{b,j}),E_{a,i}]$ is contained in $\mathfrak{n}_{\mathbf{a}}$, it should be zero and thus both $[\tilde{p}(E_{a,i}),E_{b,j}]$ and $[\tilde{p}(E_{b,j}),E_{a,i}]$ should be zero for $a\neq b$. Here we use the condition that $p>1$. So $\tilde{p}(E_{b,j})$ is zero. □

3.3. **Sub-Grassmannians.** To extend the Proposition 3.5 to the general case, we consider the following problem: Suppose that a Schubert variety $\sigma_{\mathbf{a}}$ is contained in a proper sub-Grassmannian of $Gr(m,n)$ and $\sigma_{\mathbf{b}}$ is the minimal sub-Grassmannian among them. Then will any integral variety of $\mathcal{B}_{\mathbf{a}}$ be contained in a sub-Grassmannian $\sigma_{\mathbf{b}}$?

Proposition 3.8. *Let* $\mathbf{a}=(p_1^{q_1},\cdots,p_r^{q_r})\in P(m,n)$ *be a partition and let* $\mathbf{a}'=({p'_1}^{q'_1},\cdots,{p'_r}^{q'_r})$ *be the conjugate of* $\mathbf{a}$. *Suppose that* $\mathfrak{n}_{\mathbf{a}}$ *is contained in a proper rectangle in* $\mathfrak{m}$. *Let* $\mathbf{b}$ *be the partition corresponding to the minimal rectangle among them. Then any integral variety of* $\mathcal{B}_{\mathbf{a}}$ *is contained in a sub-Grassmannian* $\sigma_{\mathbf{b}}$ *except when* $q_1+\cdots+q_r=m$ *and* $q_r=1$ *or* $q'_1+\cdots+q'_r=n-m$ *and* $q'_r=1$.

If both $q_1+\cdots+q_r<m$ and $q'_1+\cdots+q'_r<n-m$ hold, then there is no proper sub-Grassmnnian containing $\sigma_{\mathbf{a}}$. So the cases we will consider below is either when $q_1+\cdots+q_r=m$ and $q_r\geq 2$ or when $q'_1+\cdots+q'_r=n-m$ and $q'_r\geq 2$.

Lemma 3.9. *Let* $\mathbf{a}$ *and* $\mathbf{b}$ *be two partitions such that* $\mathfrak{n}_{\mathbf{a}}$ *is a subspace of* $\mathfrak{n}_{\mathbf{b}}$ *with* $\mathcal{S}_{\mathbf{a}}\subset\mathcal{S}_{\mathbf{b}}$ *and thus there is a projection* $\mathcal{B}_{\mathbf{a}}\to\mathcal{B}_{\mathbf{b}}$ *as in proposition 3.2. Assume that* $\mathcal{B}_{\mathbf{b}}$ *is strongly rigid. For* $p:\mathfrak{n}_{\mathbf{a}}\to\mathfrak{m}_{\mathbf{a}}$, *define* $\tilde{p}$ *by the composite map* $\mathfrak{n}_{\mathbf{a}}\to\mathfrak{m}_{\mathbf{a}}\twoheadrightarrow\mathfrak{m}_{\mathbf{b}}$. *If* $\tilde{p}=0$ *for all* $p\in H^{1,1}(B_{\mathbf{a}})$, *then any integral subvariety of* $\mathcal{B}_{\mathbf{a}}$ *is contained in an integral variety of* $\mathcal{B}_{\mathbf{b}}$, *which is a Schubert variety of type* $\mathbf{b}$.

Proof. The proof is similar to the proof of Lemma 3.6. □

Proof of Proposition 3.8 First, we consider the case $\mathbf{a} = (p_1^{q_1}, p_2^{q_2}, p_3^{q_3})$, $p_3 \neq 0$ is a partition with $q_1 + q_2 + q_3 = m$ and $q_3 \geq 2$. As the proof will show, the general case can be obtained in the same way.

Claim. Put $\mathbf{b} = (p_3^m)$.

$$\mathbf{a} = \begin{array}{|c|c|c||c|c|c|}\hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & \bullet & 0 & 0 & 0 & 0 \\ \hline\hline 0 & * & * & 0 & 0 & 0 \\ \hline 0 & 0 & * & \bullet & 0 & 0 \\ \hline 0 & 0 & 0 & \bullet & \bullet & 0 \\ \hline \end{array} \qquad \mathbf{b} = \begin{array}{|c|c|c||c|c|c|}\hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline\hline * & * & * & 0 & 0 & 0 \\ \hline * & * & * & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & \bullet & \bullet & 0 \\ \hline \end{array}$$

Then any integral variety of $\mathcal{B}_{\mathbf{a}}$ is contained in a Schubert variety of type $\mathbf{b}$.

Proof. Let $p : \mathfrak{n}_{\mathbf{a}} \to \mathfrak{m}_{\mathbf{a}}$ be a map with $\partial p = 0$ Then $[p(X), Y] - [p(Y), X] \in \mathfrak{n}_{\mathbf{a}}$ for all $X, Y \in \mathfrak{n}_{\mathbf{a}}$. Put $\tilde{p} : \mathfrak{n}_{\mathbf{a}} \to \mathfrak{m}_{\mathbf{b}}$ to be the composition of p with the projection $\mathfrak{m}_{\mathbf{a}} \to \mathfrak{m}_{\mathbf{b}}$.

Since $[\mathfrak{m}_{\mathbf{a}}/\mathfrak{m}_{\mathbf{b}}, \mathfrak{n}_{\mathbf{a}}] \subset \mathfrak{n}_{\mathbf{b}}$, we have $[\tilde{p}(X), Y] - [\tilde{p}(Y), X] \in \mathfrak{n}_{\mathbf{b}}$ for all $X, Y \in \mathfrak{n}_{\mathbf{a}}$. Assume that $q_1 + 1 \leq k, \ell \leq q_1 + q_2$, $q_1 + q_2 + 1 \leq i, j \leq m$, $m + 1 \leq r, s \leq n - p_1$ and $n - p_1 + 1 \leq a, b \leq n - p_2$. Note that for $X \in \mathfrak{m}_{\mathbf{b}}$, if $[X, E_{r,i}] = [X, E_{a,i}] = 0$ for all r and a for a fixed i, then X=0.

$$\begin{array}{|c|c|c||c|c|c|}\hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & \bullet & 0 & 0 & 0 & 0 \\ \hline\hline 0 & E_{r,k} & E_{r,i} & 0 & 0 & 0 \\ \hline 0 & 0 & E_{a,i} & \bullet & 0 & 0 \\ \hline 0 & 0 & 0 & \mathfrak{m}_{\mathbf{b}} & \mathfrak{m}_{\mathbf{b}} & 0 \\ \hline \end{array}$$

From $[\tilde{p}(E_{r,k}), E_{s,i}] - [\tilde{p}(E_{s,i}), E_{r,k}] \in \mathfrak{n}_{\mathbf{b}}$, we see that both $[\tilde{p}(E_{r,k}), E_{s,i}]$ and $[\tilde{p}(E_{s,i}), E_{r,k}]$ are contained in $\mathfrak{n}_{\mathbf{b}}$. So $[\tilde{p}(E_{r,k}), E_{s,i}] = 0$. The same equation hold if we replace $E_{s,i}$ by $E_{a,i}$. Thus $\tilde{p}(E_{r,k}) = 0$.

Put $\mathbf{c} = ((n-m)^{(q_1+q_2)}, p_3^{q_3})$.

$$\mathbf{c} = \begin{array}{|c|c|c||c|c|c|}\hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & \bullet & 0 & 0 & 0 & 0 \\ \hline\hline 0 & 0 & * & 0 & 0 & 0 \\ \hline 0 & 0 & * & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & \bullet & \bullet & 0 \\ \hline \end{array}$$

Then $\tilde{p}$ restricts to a map $\mathfrak{n}_{\mathbf{c}} \to \mathfrak{m}_{\mathbf{b}} \subset \mathfrak{m}_{\mathbf{c}}$. Also we have $[\tilde{p}(X), Y] - [\tilde{p}(Y), X] \in \mathfrak{n}_{\mathbf{c}}$ for all $X, Y \in \mathfrak{n}_{\mathbf{c}}$. Since $q_3 \geq 2$, $\mathcal{B}_{\mathbf{c}}$ is strongly rigid so that $\tilde{p}$

is zero on $\mathfrak{n}_{\mathbf{c}}$. Hence $\tilde{p}$ is zero. By Lemma 3.9, any integral variety of $\mathcal{B}_{\mathbf{a}}$ is contained in a Schubert variety of type $\mathbf{b}$.

In the same way, we can show that any integral manifold of $\mathcal{B}_{\mathbf{a}}$ is contained in a Schubert variety of type $\hat{\mathbf{b}}$ for $\hat{\mathbf{b}} = (p_3'^{\,n-m})$.

$$\mathbf{a} = \begin{array}{|c|c|c||c|c|c|}\hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & \bullet & 0 & 0 & 0 & 0 \\ \hline\hline 0 & * & * & 0 & 0 & 0 \\ \hline 0 & 0 & * & \bullet & 0 & 0 \\ \hline 0 & 0 & 0 & \bullet & \bullet & 0 \\ \hline \end{array} \qquad \hat{\mathbf{b}} = \begin{array}{|c|c|c||c|c|c|}\hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & 0 & 0 & 0 & 0 & 0 \\ \hline\hline 0 & * & * & 0 & 0 & 0 \\ \hline 0 & * & * & 0 & 0 & 0 \\ \hline 0 & * & * & 0 & 0 & 0 \\ \hline \end{array}$$

Here we use the rigidity of the Schubert differential system $\mathcal{B}_{\hat{\mathbf{c}}}$ with $\hat{\mathbf{c}} = (m^{(q_1'+q_2')}, p_3'^{\,q_3'}), q_3' \geq 2$.

$$\hat{\mathbf{c}} = \begin{array}{|c|c|c||c|c|c|}\hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & 0 & 0 & 0 & 0 & 0 \\ \hline \bullet & 0 & 0 & 0 & 0 & 0 \\ \hline\hline 0 & * & * & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & \bullet & 0 & 0 \\ \hline 0 & 0 & 0 & \bullet & 0 & 0 \\ \hline \end{array}$$

□

Theorem 3.10. *Let* $\mathbf{a} = (p_1^{q_1}, \cdots, p_r^{q_r})$ *be a partition and let* $\mathbf{a}' = (p_1'^{\,q_1'}, \cdots, p_r'^{\,q_r'})$ *be its conjugate. Then* $\sigma_{\mathbf{a}}$ *is Schubert rigid if* $q_i, q_i' \geq 2$ *for all* $i \leq r$.

Proof. We will use the induction on r. Thanks to Proposition 3.8, we may assume that there is no proper sub-Gassmannian containing $\sigma_{\mathbf{a}}$ so that $q_1 + \cdots q_r < m$ and $q_1' + \cdots q_r' < n - m$.

By Lemma 3.6 and the same argument as in Proposition 3.5, any integral varieties of $\mathcal{B}_{\mathbf{a}}$ is foliated by the Schubert varieties of type $\mathbf{b} = ((n-m)^q)$, where $q = q_1 + \cdots q_r$. Then the space of leaves A of this foliation will be an integral variety of the Schubert differential system $\mathcal{B}_{\mathbf{c}}$ on $Gr(q,n)$, $\mathbf{c} = ((p_1 + (m-q))^{q_1}, \cdots, (p_r + (m-q))^{q_r})$. By Proposition 3.8, any integral varieties of $\mathcal{B}_{\mathbf{c}}$ is contained in a sub-Grassmannian of type $((p_r + (m-q))^q)$ in $Gr(q,n)$. Thus A is an integral varieties of the Schubert differential system $\mathcal{B}_{\mathbf{d}}$, $\mathbf{d} = ((p_1 - p_r)^{q_1}, \cdots, (p_{r-1} - p_r)^{q_{r-1}})$ on this sub-Grassmannian, which is rigid by the induction hypothesis. □

4. SCHUR RIGIDITY

4.1. **Criterions for the equality $B_{\mathbf{a}} = R_{\mathbf{a}^*}$.** Let $\sigma_{\mathbf{a}}$ be a Schubert variety. If $\sigma_{\mathbf{a}}$ is Schubert rigid and the Schur differential system $\mathcal{R}_{\mathbf{a}^*}$ is equal to the Schubert differential system $\mathcal{B}_{\mathbf{a}}$, then $\sigma_{\mathbf{a}}$ is Schur rigid. To compare $\mathcal{B}_{\mathbf{a}}$ and $\mathcal{R}_{\mathbf{a}^*}$, we will use the following description of $\mathcal{R}_{\mathbf{a}^*}$.

Proposition 4.1. *Let* $B_{\mathbf{a}}$ *(resp.* $R_{\mathbf{a}^*}$*) be the fiber of* $\mathcal{B}_{\mathbf{a}}$ *(resp.* $\mathcal{R}_{\mathbf{a}^*}$*) at* $E \in Gr(m,n)$. *Then* $R_{\mathbf{a}^*}$ *is equal to*

$$Gr(|\mathbf{a}^*|, E^* \otimes Q) \cap \mathbb{P}(\mathbb{S}_{\mathbf{a}^*}(E^*) \otimes \mathbb{S}_{(\mathbf{a}^*)'}(Q)) \subset \mathbb{P}(\wedge^{|\mathbf{a}^*|}(E^* \otimes Q))$$

and $B_{\mathbf{a}}$ *is the orbit of a highest weight vector in the irreducible representation space* $\mathbb{P}(\mathbb{S}_{\mathbf{a}^*}(E^*) \otimes \mathbb{S}_{(\mathbf{a}^*)'}(Q))$ *of* $SL(E^*) \times SL(Q)$.

Proof. By Proposition 3.1 of [Ho]. □

Note that we adapt the convention that $\sigma_{\mathbf{a}}$ is a Schubert variety of dimension $|\mathbf{a}^*|$, while X_w is a Schubert variety of dimension $\ell(w)$ in [Ho]. Thus, when $\mathbf{a}^* \in P(m,n)$ corresponds to $w \in W^P$, $\mathcal{B}_{\mathbf{a}}$ is equal to $\mathcal{B}_w$ and $\mathcal{R}_{\mathbf{a}^*}$ is equal to $\mathcal{R}_w$.

Since P_0 acts on the fiber $B_{\mathbf{a}}$, we have a decomposition $\mathfrak{p}_0 = \mathfrak{m}_{\mathbf{a}} + \mathfrak{l}_{\mathbf{a}} + \mathfrak{m}^*_{\mathbf{a}}$, where the tangent space of $B_{\mathbf{a}}$ at a point is isomorphic to $\mathfrak{m}_{\mathbf{a}}$. A sufficient condition for the equality $B_{\mathbf{a}} = R_{\mathbf{a}^*}$ is given by Proposition 3.2 in [Ho]. Put $\mathbf{I}_{\mathbf{a}} = \mathbb{S}_{\mathbf{a}^*}(E^*) \otimes \mathbb{S}_{(\mathbf{a}^*)'}(Q)$.

Proposition 4.2. *Assume that for the* $\mathfrak{l}_{\mathbf{a}}$*-generators* φ *of all irreducible* $\mathfrak{l}_{\mathbf{a}}$*-representation spaces in the complement of* $\mathfrak{m}_{\mathbf{a}}$ *in* $\mathfrak{n}^*_{\mathbf{a}} \otimes \mathfrak{m}/\mathfrak{n}_{\mathbf{a}}$, *we have*

$$\varphi^k(v_1 \wedge \cdots \wedge v_k) \notin \mathbf{I}_{\mathbf{a}},$$

where $\{v_1, \cdots, v_k\}$ *is a basis of* $\mathfrak{n}_{\mathbf{a}}$ *and* $\varphi^k : \wedge^k \mathfrak{n}_{\mathbf{a}} \to \wedge^k \mathfrak{m} / \wedge^k \mathfrak{n}_{\mathbf{a}}$ *is defined by*

$$\varphi^k(v_1 \wedge \cdots \wedge v_k) = \sum_i v_1 \wedge \cdots \wedge \varphi(v_i) \wedge \cdots \wedge v_k \mod \wedge^k \mathfrak{n}_{\mathbf{a}}.$$

Then $B_{\mathbf{a}}$ *is equal to* $R_{\mathbf{a}^*}$.

Let E be an m-subspace of $\mathbb{C}^n$ and Q be the quotient $\mathbb{C}^n/E$. Then $\mathfrak{m}$ is equal to $E^* \otimes Q$. Take a partition $\mathbf{a} = (p_1^{q_1}, \cdots, p_r^{q_r}) \in P(m,n)$ and let $\mathbf{a}' = ({p'_1}^{q'_1}, \cdots, {p'_r}^{q'_r})$, $p'_r \neq 0$ be its conjugate. Write $E = \oplus_{i=1}^{r_E} E_i$ and $Q = \oplus_{a=1}^{r_Q} Q_a$ so that $\mathfrak{l}_{\mathbf{a}} = (\oplus_{i=1}^{r_E} sl(E_i)) \oplus (\oplus_{a=1}^{r_Q} sl(Q_a))$. Note that r_E is r if $q_1 + \cdots + q_r = m$ and is $r+1$, otherwise, and r_Q is r if $q'_1 + \cdots q'_r = n-m$ and is $r+1$, otherwise. We indexed E_i and Q_a keeping the order of the basis $\{e_1, \cdots, e_n\}$ of $\mathbb{C}^n$ such that $\{e_1, \cdots, e_m\}$ is a basis of E. Set $r_i = \dim E_i$ and $s_a = \dim Q_a$ for $1 \leq i \leq r_E$ and $1 \leq a \leq r_Q$. Then $E_i =< e_{r_{i-1}+1}, \cdots, e_{r_i} >$ and $Q_a =< q_{s_{a-1}+1}, \cdots, q_{s_a} >$, where $q_p := e_{m+p}$ for $p = 1, \cdots, n-m$.

Let Π be the index set of (i,a) such that $\mathfrak{n}_{\mathbf{a}} = \oplus_{(i,a)\in\Pi}(E_i^* \otimes Q_a)$. As a subspace of $\mathfrak{n}^*_{\mathbf{a}} \otimes \mathfrak{m}/\mathfrak{n}_{\mathbf{a}}$, $\mathfrak{m}_{\mathbf{a}} = (\oplus_{i<j} E_i^* \otimes E_j) \oplus (\oplus_{b<a} Q_b^* \otimes Q_a)$ is equal to

$$(\oplus_{i<j} E_i^* \otimes E_j \otimes < Id_{\oplus_{a\in\Pi_{i,j}} Q_a} >_{\mathbb{C}}) \oplus (\oplus_{b<a} < Id_{\oplus_{i\in\Pi_{b,a}} E_i} >_{\mathbb{C}} \otimes Q_b^* \otimes Q_a)$$

where $\Pi_{i,j} = \{a : (i,a) \notin \Pi, (j,a) \in \Pi\}$ and $\Pi_{b,a} = \{i : (i,b) \in \Pi, (i,a) \notin \Pi\}$.

We may choose the order of the set of roots of $SL(n)$ in such a way that the maximal root is located in the most left and the lowest box $E_1^* \otimes Q_s$ and the minimal root is located in the most right and the highest box $E_r^* \otimes Q_1$. Then the highest weight vector in $\mathfrak{m}_{\mathbf{a}} \subset \mathfrak{n}_{\mathbf{a}}^* \otimes \mathfrak{m}/\mathfrak{n}_{\mathbf{a}}$ is either

$\sum_{a \in \Pi_{i,j}} \sum_{q_p \in Q_a} x_{\alpha_p}^* \otimes x_{\beta_p}$, where $x_{\alpha_p} = e_{r_j}^* \otimes q_p$ and $x_{\beta_p} = e_{r_{i-1}+1}^* \otimes q_p$ for some $i < j$, or,

$\sum_{i \in \Pi_{b,a}} \sum_{e_p \in E_i} x_{\alpha_p}^* \otimes x_{\beta_p}$, where $x_{\alpha_p} = e_p^* \otimes q_{s_{b-1}+1}$ and $x_{\beta_p} = e_p^* \otimes q_{s_a}$ for some $b < a$,

For example, Consider $\mathbf{a} = (9^2, 7^2, 3^4)$ in $P(10, 19)$. Then we have $E = \oplus_{i=1}^4 E_i$, $Q = \oplus_{a=1}^3 Q_a$ and $\Pi = \{(2,1), (3,1), (3,2), (4,1), (4,2), (4,3)\}$. The highest weight vector of $\mathfrak{m}_{\mathbf{a}}$ is



Proposition 4.3. *Let $E = \oplus_i E_i$ and $Q = \oplus_a Q_a$ be the decomposition associate to $\mathbf{a}$ as in the above and $\{e_1, \cdots, e_n\}$ be a basis for $\mathbb{C}^n$ indexed as in the above. Then the highest weight vector of an irreducible $\mathfrak{l}_{\mathbf{a}}$-representation space in the complement of $\mathfrak{m}_{\mathbf{a}}$ in $\mathfrak{n}_{\mathbf{a}}^* \otimes \mathfrak{m}/\mathfrak{n}_{\mathbf{a}}$ is either*

(1) a decomposable vector $x_\alpha^* \otimes x_\beta$, where $x_\alpha = e_{r_j}^* \otimes q_{s_{b-1}+1}$ is the lowest weight vector of $E_j^* \otimes Q_b$ and $x_\beta = e_{r_{i-1}+1}^* \otimes q_{s_a}$ is the highest weight vector of $E_i^* \otimes Q_a$ for some $(j, b) \in \Pi, (i, a) \notin \Pi$, or,

(2) $\sum_{q_p \in Q_a} x_{\alpha_p}^* \otimes x_{\beta_p}$, where $x_{\alpha_p} = e_{r_j}^* \otimes q_p$ and $x_{\beta_p} = e_{r_{i-1}+1}^* \otimes q_p$ for some $(j, a) \in \Pi, (i, a) \notin \Pi$ such that $(j, a-1) \in \Pi$ and $(i, a-1) \notin \Pi$, or,

(3) $\sum_{e_p \in E_i} x_{\alpha_p}^* \otimes x_{\beta_p}$, where $x_{\alpha_p} = e_p^* \otimes q_{s_{b-1}+1}$ and $x_{\beta_p} = e_p^* \otimes q_{s_a}$ for some $(i, b) \in \Pi, (i, a) \notin \Pi$ such that $(i+1, b) \in \Pi$ and $(i+1, a) \notin \Pi$.

Proof. If $i \neq j$ and $a \neq b$, then $(E_j^* \otimes Q_b)^* \otimes (E_i^* \otimes Q_a)$ is an irreducible $\mathfrak{l}_{\mathbf{a}}$-representation space. But if $i \neq j$ and $a = b$, then $(E_j^* \otimes Q_a)^* \otimes (E_i^* \otimes Q_a) \simeq E_j \otimes E_i^* \otimes (Q_a^* \otimes Q_a)$ is decomposed as $(E_j \otimes E_i^* \otimes < Id_{Q_a} >_{\mathbb{C}}) \oplus (E_j \otimes E_i^* \otimes sl(Q_a))$, each of which are irreducible $sl(E_j) \times sl(E_i) \times sl(Q_a)$-representation spaces. The highest weight vector of the irreducible representation spaces $(E_j^* \otimes Q_b)^* \otimes (E_i^* \otimes Q_a)$ or $(E_j \otimes E_i^* \otimes sl(Q_a))$ is of type (1).

The component $E_i^* \otimes E_j$ of $\mathfrak{m}_{\mathbf{a}}$ corresponds to the component $E_i^* \otimes E_j \otimes < Id_{\oplus_{a \in \Pi_{i,j}} Q_a} >_{\mathbb{C}}$ in $\mathfrak{n}_{\mathbf{a}}^* \otimes \mathfrak{m}/\mathfrak{n}_{\mathbf{a}}$, so its complement in $\oplus_{a \in \Pi_{i,j}} E_i^* \otimes E_j \otimes < Id_{Q_a} >_{\mathbb{C}}$ is $\oplus_{a \in \tilde{\Pi}_{i,j}} E_i^* \otimes E_j \otimes < Id_{Q_a} >_{\mathbb{C}}$, where $\tilde{\Pi}_{i,j} = \{a : (i, a) \notin \Pi, (j, a) \in$

$\Pi, (i, a-1) \notin \Pi, (j, a-1) \in \Pi\}$ is obtained from $\Pi_{i,j}$ by excluding the smallest index in $\Pi_{i,j}$. The highest weight of the irreducible representation space in these components is of type (2).

Considering the case when $i = j$ and $a \neq b$, we obtain the highest weight vectors of type (3). □

For example, the highest weight vector $\sum_{q_p \in Q_a} x^*_{\alpha_p} \otimes x_{\beta_p}$ of type (2) is



E_1 E_2 E_3 E_4
Q_1
Q_2
Q_3
$\times_3$ $\times_4$ $\times_5$ $\times_6$
$\bullet_3$ $\bullet_4$ $\bullet_5$ $\bullet_6$
$x_{\beta_p} = \times_p$ $x_{\alpha_p} = \bullet_p$

or



E_1 E_2 E_3 E_4
Q_1
Q_2
Q_3
$\times_7$ $\times_8$ $\times_9$
$\bullet_7$ $\bullet_8$ $\bullet_9$
$x_{\beta_p} = \times_p$ $x_{\alpha_p} = \bullet_p$

etc.

4.2. **Proof of the equality $B_{\mathbf{a}} = R_{\mathbf{a}^*}$.** We will use the same notations as in the previous section. When $\{v_1, \cdots, v_k\}$ is a basis of $\mathfrak{n}_{\mathbf{a}}$ such that $v_1 = x_\alpha$, we will use the notation $x_\beta \wedge \hat{x}_\alpha \wedge \cdots \wedge v_k$ to denote the k-vector obtained from $v_1 \wedge \cdots \wedge v_k$, which may be considered as a base k-vector, by replacing $v_1 = x_\alpha$ with x_β.

We call the union of all the columns corresponding to e_p's in E_i the E_i-*column.* Similarly, we call the union of all the rows corresponding e_p's in Q_a the Q_a-*row.*

Proposition 4.4. *Let* $\mathbf{a} = (p_1^{q_1}, \cdots, p_r^{q_r})$ *be a partition and let* $\mathbf{a}' = ({p'_1}^{q'_1}, \cdots, {p'_r}^{q'_r})$ *be its conjugate. Take a decomposition* $E = \oplus_i E_i$ *and* $Q = \oplus_a Q_a$ *associated to* $\mathbf{a}$ *as in the previous section. Let* Π *be the index set of* (i, a) *such that* $\mathfrak{n}_{\mathbf{a}} = \oplus_{(i,a)\in\Pi} E_i^* \otimes Q_a$. *Then the Schubert differential system* $\mathcal{B}_{\mathbf{a}^*}$ *is equal to the Schur differential system* $\mathcal{R}_{\mathbf{a}}$ *except when both* E_i *and* Q_a *are one dimensional for some* $(i, a) \notin \Pi$ *with* $E_i^* \otimes Q_a$ *adjacent to* $\mathfrak{n}_{\mathbf{a}}$.

Proof. We will divided the proof by two: I. when the highest weight of the irreducible component in the complement of $\mathfrak{m}_{\mathbf{a}}$ in $\mathfrak{n}^*_{\mathbf{a}} \otimes \mathfrak{m}/\mathfrak{n}_{\mathbf{a}}$ is of type (1) and II. when it is of type (2) or (3) (See Proposition 4.3 for the types of the highest weight vectors).

I. Type (1): Fix $(j, b), \in \Pi, (i, a) \notin \Pi$. Let $\{v_1, \cdots, \cdots, v_k\}$ be a basis of $\mathfrak{n}_{\mathbf{a}}$ with $v_1 = x_\alpha$ is a lowest weight vector of $E_j^* \otimes Q_b$ and x_β is the highest weight vector of $E_i^* \otimes Q_a$. We will show that $x_\beta \wedge \hat{x}_\alpha \wedge \cdots \wedge v_k$ has a nonzero component in $\mathbf{I_b}$ for a partition $\mathbf{b} \neq \mathbf{a}$ with $|\mathbf{b}| = |\mathbf{a}|$.

Case 1. Assume that $q_i' \geq 2$ for all i. Then we can take $\mathbf{b}$ with $\mathfrak{n}_{\mathbf{b}} = \mathfrak{n}_{\mathbf{a}} - \{x_{\beta'}\} \cup \{x_{\alpha'}\}$ where $x_{\beta'}$ is the highest weight in the boxes $E_j^* \otimes Q_c$ for all c with $(j, c) \in \Pi$ and $x_{\alpha'}$ is the lowest weight vector in the boxes $E_k^* \otimes Q_a$ for all k with $(k, a) \notin \Pi$. Note that there is $x_{\gamma_j}, j = 1, \cdots, 4$ in $\mathfrak{p}_0$ such that $ad(x_{\gamma_2})ad(x_{\gamma_1})x_\beta = x_{\alpha'}$ and $ad(x_{\gamma_4})ad(x_{\gamma_3})x_{\beta'} = x_\alpha$ and $ad(x_{\gamma_j}), j = 1, \cdots, 4$

If x_α, x_β, $x_{\beta'}$ and $x_{\alpha'}$ lie neither in the same E_i-column nor in the same Q_a-raw as in the picture,


$\mathfrak{n}_{\mathbf{a}}$
$\mathfrak{n}_{\mathbf{b}}$
$\xrightarrow{ad(x_{\gamma_4})}$
$ad(x_{\gamma_3})$ ↑
$ad(x_{\gamma_1})$ ↑
$\xrightarrow{ad(x_{\gamma_2})}$
⟶

$x_\alpha =$ one of •'s

$x_\beta =$ one of ×'s

$x_{\beta'} = \circ$

$x_{\alpha'} = \star$

then all γ_i are distinct. Thus

$$ad(x_{\gamma_4}) \cdots ad(x_{\gamma_1})(x_\beta \wedge \cdots \wedge \hat{x}_\alpha \wedge \cdots \wedge v_k) = x_{\alpha'} \wedge \cdots \wedge \hat{x}_{\beta'} \wedge \cdots \wedge v_k.$$

But $x_{\alpha'} \wedge \cdots \wedge \hat{x}_{\beta'} \wedge \cdots \wedge v_k$ is the lowest weight vector of $\mathbf{I_b}$. Since $ad(\mathfrak{p}_0)$ preserves $\mathbf{I_b}$, there is a nonzero $\mathbf{I_b}$-component in $x_\beta \wedge \cdots \wedge \hat{x}_\alpha \wedge \cdots \wedge v_k$ and thus it is not contained in $\mathbf{I_a}$.

If x_α, x_β, $x_{\beta'}$ and $x_{\alpha'}$ lie either in the same E_i-column or in the same Q_a-row as in the picture



then $\gamma_2 = \gamma_4$ or $\gamma_1 = \gamma_3$. We consider the case when $\gamma_2 = \gamma_4$(the proof for the other case is similar to this case). Thus the multivector $x_{\alpha'} \wedge \hat{x}_{\beta'} \wedge \cdots \wedge v_k \in \mathbf{I_b}$ can be obtained from the multivector $x_\beta \wedge \hat{x}_\alpha \wedge \cdots \wedge v_k$ by applying $g \circ ad(x_{\gamma_3})ad(x_{\gamma_1})$, where $g \in SL(m) \times SL(n-m)$ which exchanges the most left column $e^*_{s_{i-1}+1} \otimes q_p\,, 1 \leq p \leq n-m$ with the most right column $e^*_{s_i} \otimes q_p\,, 1 \leq p \leq n-m$ in the E_i-column. This shows that $x_\beta \wedge \hat{x}_\alpha \wedge \cdots \wedge v_k$ is not contained in $\mathbf{I_a}$.

Case 2. If some $q'_a = 1$, then there may be no partition $\mathbf{b}$ with such property as in Case 1. This is the case when dim Q_a is one and $a = b+1$ and $E^*_{j+1} \otimes Q_a \notin \mathfrak{n}_{\mathbf{a}}$. Then $x_{\alpha'}$ is left to $x_{\beta'}$ and they are adjacent so we cannot find such a partition $\mathbf{b}$.

$\mathfrak{n}_{\mathbf{a}}$



$\mathfrak{n}_{\mathbf{b}}$



But in this case, consider $\mathbf{b}'$ such that $\mathfrak{n}_{\mathbf{b}'} = \mathfrak{n}_{\mathbf{a}} - \{x_{\beta''}\} \cup \{x_{\alpha'}\}$ where $x_{\beta''}$ is the highest weight vector in the boxes $E_k^* \otimes Q_b$ for all k with $(k, b) \in \Pi$ and $x_{\alpha'}$ is the lowest weight vector in the boxes $E_i^* \otimes Q_c$ for all c with $(i, c) \notin \Pi$.

$\mathfrak{n}_{\mathbf{b}'}$



If $i \neq j+1$, then there is such a partition $\mathbf{b}'$. If $i = j+1$, then, by the assumption, dim E_i is not equal to one and thus we can find such a partition $\mathbf{b}'$. Then by the same argument as in the case 1, we can prove that $x_\beta \wedge \hat{x}_\alpha \wedge \cdots \wedge v_k$ is not contained in $\mathbf{I}_{\mathbf{a}}$.

II. Type(2) or (3) Let $\sum_{q_p \in Q_a} x^*_{\alpha_p} \otimes x_{\beta_p}$ be the highest weight vector of type (2) of an irreducible representation space in the complement of $\mathfrak{m}_{\mathbf{a}}$ in $\mathfrak{n}^*_{\mathbf{a}} \otimes \mathfrak{m}/\mathfrak{n}_{\mathbf{a}}$, where $x_{\alpha_p} = e^*_{r_j} \otimes q_p$ and $x_{\beta_p} = e^*_{r_{i-1}+1} \otimes q_p$ for some $(j, a) \in \Pi, (i, a) \notin \Pi$ such that $(j, a-1) \in \Pi$ and $(i, a-1) \notin \Pi$. We will show that

$\sum_{q_p\in Q_a} x_{\beta_p}\wedge\hat{x}_{\alpha_p}\wedge\cdots\wedge v_k$ has a nonzero component in $\mathbf{I_b}+\mathbf{I_c}$ if $\dim Q_a\geq 2$, and has a nonzero component in $\mathbf{I_b}$, otherwise, for some partition $\mathbf{b}$ and $\mathbf{c}$.

Applying the adjoint actions successively to

$$\sum_{q_p\in Q_a} x_{\beta_p}\wedge\hat{x}_{\alpha_p}\wedge\cdots\wedge v_k = x_{\beta_{s_{a-1}+1}}\wedge\hat{x}_{\alpha_{s_{a-1}+1}}\wedge\cdots\wedge v_k+\sum_{p=s_{a-1}+2}^{p=s_a} x_{\beta_p}\wedge\hat{x}_{\alpha_p}\wedge\cdots\wedge v_k,$$

we can get

$$x_{\beta_{s_{a-1}+1}}\wedge\hat{x}_{\alpha_{s_c}}\wedge\cdots\wedge v_k \ + \ x_{\beta_{s_{a-2}+1}}\wedge\hat{x}_{\alpha_{s_c}}\wedge\cdots\wedge v_k,$$

where $x_{\beta_{s_{a-2}+1}} = e^*_{r_{i-1}+1}\otimes q_{s_{a-2}+1}$ and $x_{\alpha_{s_c}} = e^*_{r_j}\otimes q_{\alpha_{s_c}}$ and c is the largest index c such that $(j,c)\in\Pi$.



$x_{\beta_p} = \times_p \quad x_{\alpha_p} = \bullet_p$

Applying the adjoint actions again, we can get

$$y_{\beta_{s_{a-1}+1}}\wedge\hat{y}_{\alpha_{s_c}}\wedge\cdots\wedge v_k \ + \ y_{\beta_{s_{a-2}+1}}\wedge\hat{y}_{\alpha_{s_c}}\wedge\cdots\wedge v_k,$$

where $y_{\beta_{s_{a-1}+1}} = e^*_{r_h}\otimes q_{s_{a-1}+1}$ and $y_{\beta_{s_{a-2}+1}} = e^*_{r_i}\otimes q_{s_{a-2}+1}$ and $y_{\alpha_{s_c}} = e^*_{r_{j-1}+1}\otimes q_{\alpha_{s_c}}$ and h is the largest index h such that $(h,a)\notin\Pi$. This is the sum of the lowest weight vector of $\mathbf{I_b}$ and that of $\mathbf{I_c}$ for some partition $\mathbf{b}$ and $\mathbf{c}$ such that all $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$ are distinct.



□

By Theorem 3.10 and Proposition 4.4, we get

Theorem 4.5. *Let* $\mathbf{a} = (p_1^{q_1}, \cdots, p_r^{q_r}), p_r \neq 0$ *be a partition and let* $\mathbf{a}' = ({p'_1}^{q'_1}, \cdots, {p'_r}^{q'_r})$, $p'_r \neq 0$ *be its conjugate. Then* $\sigma_{\mathbf{a}}$ *is Schur rigid if* $q_i, q'_i \geq 2$ *for all* $i \leq r$.

Remark 4.6. One of the problems in algebraic geometry is the smoothability of a singular Schubert variety X_w of G/P. We say X_w is *smoothable* if there is a smooth subvariety X of G/P with $[X] = [X_w]$ in $H_*(G/P, \mathbb{Z})([\mathrm{B}])$. Assume that $\mathbf{a} = (p^q)$ and that $p = 1$ or $q = 1$ but both are not 1. Then $\sigma_{\mathbf{a}}$ is a singular Schubert variety and the non-smoothability of $X_{\mathbf{a}}$ is proved in [B]: if X is a subvariety of $Gr(m, n)$ with $[X] = [\sigma_{\mathbf{a}}]$, then X is a Schubert variety of type $\mathbf{a}$. By Theorem 4.5, for a partition $\mathbf{a} = (p_1^{q_1}, \cdots, p_r^{q_r})$ with its conjugate $\mathbf{a}' = ({p'_1}^{q'_1}, \cdots, {p'_r}^{q'_r})$, if $q_i, q'_i \geq 2$, for all i, then the singular Schubert varieties $\sigma_{\mathbf{a}}$ of type $\mathbf{a}$ is not smoothable, neither.

Remark 4.7. With the same notations as in Proposition 4.4, if both E_i and Q_a are one dimensional for some $(i, a) \notin \Pi$ with $E_i^* \otimes Q_a$ adjacent to $\mathfrak{n}_{\mathbf{a}}$, then $B_{\mathbf{a}}$ is a proper subvariety of $R_{\mathbf{a}^*}$.

Consider the highest weight vector $x_\alpha^* \otimes x_\beta$ of an irreducible component in the complement of $\mathfrak{m}_{\mathbf{a}}$ in $\mathfrak{n}_{\mathbf{a}}^* \otimes \mathfrak{m}/\mathfrak{n}_{\mathbf{a}}$ such that $x_\alpha \in \mathfrak{n}_{\mathbf{a}}$ and $x_\beta \in E_i^* \otimes Q_a \subset \mathfrak{m}/\mathfrak{n}_{\mathbf{a}}$. Then one can check that $x_\beta \wedge \hat{x}_\alpha \wedge \cdots \wedge v_k$ is contained in $\mathbf{I}_{\mathbf{a}}$. Thus this gives a nontrivial element in $T_{\mathbf{a}} := T_{[\mathfrak{n}_{\mathbf{a}}]} Gr(k, \mathfrak{m}) \cap T_{\wedge^k \mathfrak{n}_{\mathbf{a}}} \mathbb{P}(\mathbf{I}_{\mathbf{a}}) \subset \mathfrak{n}_{\mathbf{a}}^* \otimes \mathfrak{m}/\mathfrak{n}_{\mathbf{a}}$. Note that as an element of $T_{[\wedge^k \mathfrak{n}_{\mathbf{a}}]} \mathbb{P}(\mathbf{I}_{\mathbf{a}})$, this tangent vector gives the map

$$x_\alpha \wedge v_2 \wedge \cdots \wedge v_k \longmapsto x_\beta \wedge v_2 \wedge \cdots \wedge v_k \in \mathbf{I}_{\mathbf{a}} / \wedge^k \mathfrak{n}_{\mathbf{a}},$$

where $\{v_1 = x_\alpha, v_2, \cdots, v_k\}$ be a basis of $\mathfrak{n}_{\mathbf{a}}$. Then $c(t) = (x_\alpha + t x_\beta) \wedge v_2 \wedge \cdots \wedge v_k$, $t \in \mathbb{C}$, is a curve in $R_{\mathbf{a}} = Gr(k, \mathfrak{m}) \cap \mathbb{P}(\mathbf{I}_{\mathbf{a}})$ whose tangent vector is $x_\alpha^* \otimes x_\beta$. But this tangent vector is not contained in $\mathfrak{m}_{\mathbf{a}}$ and thus $B_{\mathbf{a}}$ is a proper subvariety of $R_{\mathbf{a}^*}$. This is a generalization of the counterexamples considered in [W] or Example 9 of [B].

REFERENCES

[B] R. Bryant, *Rigidity and quasi-rigidity of extremal cycles in compact Hermitian symmetric spaces*, Annals of mathematics studies 153, Princeton University Press, 2004

[FH] W. Fulton and J. Harris, Representation Theory; A First Course, Springer-Verlag, 1991

[G] A. B. Goncharov, *Generalized conformal structures on manifolds*, Selecta Mathematica Sovietica, vol 6, No 4, 1987.

[H] J. Harris, Algebraic geometry, GTM 133, Springer-Verlag, 1992

[Ho] J. Hong, Rigidity of smooth Schubert varieties in Hermitian symmetric spaces, math.DG/0410138

[W] M. Walters, *Geometry and uniqueness of some extreme subvarieties in complex Grassmannians*, Ph.D. thesis, University of Michigan, 1997.

Research Institute of Mathematics

Seoul National University
San 56-1 Sinrim-dong Kwanak-gu
Seoul, 151-747 Korea
jhhong@math.snu.ac.kr